



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>A61K 31/535, 31/495, 31/50, 31/505</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 96/39145</b> <b>(43) International Publication Date:</b> 12 December 1996 (12.12.96)
<b>(21) International Application Number:</b> PCT/US96/09606 <b>(22) International Filing Date:</b> 6 June 1996 (06.06.96) <b>(30) Priority Data:</b> 08/469,147 6 June 1995 (06.06.95) US <b>(71) Applicant (for all designated States except US):</b> RHONE-POULENC RORER PHARMACEUTICALS INC. [US/US]; P.O. Box 5093, Collegeville, PA 19426-0997 (US). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> MYERS, Michael, R. [US/US]; 205 Lincoln Drive, Reading, PA 19606 (US). SPADA, Alfred, P. [US/US]; 473 Painter Way, Lansdale, PA 19446 (US). MAGUIRE, Martin, P. [US/US]; 8861 Deer Path Circle, Louisville, KY 40220 (US). PERSONS, Paul, E. [US/US]; 649 S. Henderson Road A-507, King of Prussia, PA 19406 (US). <b>(74) Agents:</b> NICHOLSON, James, A. et al.; Rhône-Poulenc Rorer Inc., P.O. Box 5093, Collegeville, PA 19426-0997 (US).		<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> PROTEIN TYROSINE KINASE ARYL AND HETEROARYL QUINAZOLINE COMPOUNDS HAVING SELECTIVE INHIBITION OF HER-2 AUTOPHOSPHORYLATION PROPERTIES  <b>(57) Abstract</b>  This invention relates to a method for the selective treatment of cell growth and differentiation characterized by activity of the human epidermal growth factor receptor type 2 (HER2). More specifically, this invention relates to the use of substituted or unsubstituted mono- or bi-cyclic aryl, heteroaryl, cycloalkyl or heterocycloalkyl compounds in selectively regulating cell growth. Pharmaceutical compositions useful for the selective treatment of cell growth and differentiation are also described.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

5

10        PROTEIN TYROSINE KINASE ARYL AND HETEROARYL QUINAZOLINE  
             COMPOUNDS HAVING SELECTIVE INHIBITION OF HER-2  
             AUTOPHOSPHORYLATION PROPERTIES

Background of the Invention

15

             This application is a continuation-in-part application of PCT International  
Application Serial No. PCT/US94/14180 having International filing date of  
December 8, 1994, designating the United States and United States Serial  
No. 08/166,199, filed December 10, 1993 which is a continuation-in-part  
20       application of United States Serial No. 07/988,515, filed December 10, 1992,  
which is a continuation-in-part application of United States Serial No.  
07/698,420, filed May 10, 1991 and PCT International Application Serial No.  
PCT/US92/03736 filed May 6, 1992 designating the United States, which  
designation has resulted in United States Patent No. 5,409,930.

25

Field of the Invention

             This invention relates to a method for the selective treatment of cell  
growth and differentiation characterized by activity of the human epidermal  
30       growth factor receptor type 2 (HER2). More specifically, this invention relates to  
the use of substituted or unsubstituted mono- or bi-cyclic aryl, heteroaryl,  
cycloalkyl or heterocycloalkyl compounds in selectively regulating cell growth.

             Normal cellular reproduction is believed to be triggered by the exposure  
35       of the cellular substrate to one or more growth factors, examples of which are  
insulin, epidermal growth factor (EGF) and platelet-derived growth factor  
(PDGF). Such growth factor receptors are imbedded in and penetrate through

the cellular membrane. The initiation of cellular reproduction is believed to occur when a growth factor binds to the corresponding receptor on the external surface of the cellular membrane. This growth factor-receptor binding alters the chemical characteristics of that portion of the receptor which exists within the cell and which functions as an enzyme to catalyze phosphorylation of either an intracellular substrate or the receptor itself, the latter being referred to as autophosphorylation. Examples of such phosphorylation enzymes include tyrosine kinases, which catalyze phosphorylation of tyrosine amino acid residues of substrate proteins.

Many disease states are characterized by the uncontrolled reproduction of cells. These disease states involve a variety of cell types and include disorders such as leukemia, cancer, psoriasis, inflammatory diseases, bone diseases, atherosclerosis and restenosis occurring subsequent to angioplastic procedures. The inhibition of tyrosine kinase is believed to have utility in the control of uncontrolled cellular reproduction, i.e., cellular proliferative disorders.

Initiation of autophosphorylation, i.e., phosphorylation of the growth factor receptor itself, and of the phosphorylation of a host of intracellular substrates are some of the biochemical events which are involved in mediator release mitogenesis and cell proliferation. Autophosphorylation of the insulin receptor and phosphorylation of substrate proteins by other receptors are the earliest identifiable biochemical hormonal responses.

Elimination of the protein tyrosine kinase (PTK) activity of the insulin receptor and of the epidermal growth factor (EGF) receptor by site-directed mutagenesis of the cellular genetic material which is responsible for generation of insulin and EGF results in the complete elimination of the receptor's biological activity. This is not particularly desirable because insulin is needed by the body to perform other biological functions which are not related to cell proliferation. Accordingly, compounds which inhibit the PTK portion of the EGF and/or PDGF receptor at concentrations less than the concentrations needed to inhibit the PTK portion of the insulin receptor could provide valuable agents for selective treatment of cell proliferation disorders.

### Reported Developments

It is generally accepted that the protein tyrosine kinase (PTK) activity associated to growth factor receptors is essential for ligand-induced biological responses, such as cell growth and differentiation. J. M. Bishop, (1985) *Cell*, **42**: 23-38 and T. Hunter (1987) *Cell*, **50**: 823-829 have reported evidence for abnormal cell growth and morphological transformation *via* tyrosine phosphorylation, growth factor receptors represent a *bonafide* oncogene family. Among these oncogene products, is the human epidermal growth factor receptor type 2 (HER2; c-erbB-2). Carpenter et al. (1987) *Annu. Rev. Biochem.*, **56**: 881- and Gill et al. (1987) *Mol. Cell. Endocrinol.*, **51**: 169- have reported that the c-erbB-1 gene encodes a 170kDa transmembrane glycoprotein. The HER2 gene is the human homolog of the rat protooncogene neu as reported by C. I. Bargmann, M. C. Hung and R. A. Weinberg (1986) *Nature*, **319**: 226-229 and its product is a 185-kDa transmembrane cell-surface glycoprotein, so called p185<sup>HER2</sup> as reported by Coussens et al. (1985) *Science*, **230**: 1132-1139 and Yamamoto et al. (1986) *Nature*, **319**: 230-234. This receptor harbors an intrinsic tyrosine-specific kinase activity which is capable of catalyzing autophosphorylation reaction as well as mediating endogenous substrate phosphorylation upon binding of their cognate ligands on the receptor extracellular domains.

The unaltered HER2 gene has been found amplified in various human tumors. In addition, a correlation between the elevated levels of p 185<sup>HER2</sup> expression and poor prognosis or short survival time has been established in patients with several cancer types. Given that the use of antibodies raised against p185<sup>HER2</sup> (that block receptor functions) results in a growth arrest of tumor-derived cell lines and anti-tumor effect in mice, strategy in which chemically-designed agents that selectively inhibit p185<sup>HER2</sup>-associated PTK are valuable in cancer therapy. It is now widely documented that numerous naturally occurring and synthetic analogues of PTK inhibitors have been systematically evaluated for their potential anti-tumor efficacy as reported by C. Workman, V. G. Brunton and D. J. Robins (1992) in *seminars in Cancer Biology*, **3**: 369-381 and C. Wasylyk, A. Gutman, R. Nicholson and B. Wasylyk (1991) *EMBO J.*, **10**: 1127-1134.

Applicants describe herein the identification and characterization of a series of quinazolin derivatives - which are tyrosine kinase inhibitors exhibiting selectivity for p185HER2. In particular, Applicants demonstrate the requirement of tyrosine kinase activity associated to HER2 gene product for maintaining transformed cell phenotype induced by overexpression of p185HER2.


### Summary of the Invention

In accordance with the present invention, there is provided a method for the selective treatment of cell growth and differentiation characterized by activity of the human epidermal growth factor receptor type 2 (HER2) comprising administering to a patient in need of such treatment an HER2 receptor inhibiting effective amount of a quinazoline compound described in this invention

Another aspect of the present invention relates to pharmaceutical compositions for selectively treating cell growth and differentiation characterized by activity of the human epidermal growth factor receptor type 2 (HER2) comprising, in admixture with a pharmaceutically acceptable carrier, a pharmaceutically effective amount of a compound of the aforementioned type.

A further aspect of this invention comprises novel compounds useful in the practice of the present method.

With respect to the aspects of this invention, the compounds described by Formula I below constitute a class of the aforementioned quinazoline compounds for use in the practice of the present invention:



Formula I

wherein

A is a substituted or unsubstituted mono- or bi-cyclic aryl, heteroaryl, cycloalkyl or heterocycloalkyl ring system of about 5 to about 12 atoms and

where the substituents may be located at any appropriate position of the ring system and are described by R;

5 X is a bond, O, S, SO, SO<sub>2</sub>, OCH<sub>2</sub>, CR<sub>4</sub>=CR<sub>4</sub>, C≡C, NR<sub>4</sub> or NR<sub>4</sub>CH<sub>2</sub>;

R independently includes hydrogen, alkyl, phenyl, halophenyl, aralkyl, hydroxy, alkoxy, aryloxy, acyloxy, halo, haloalkyl, amino, mono- and di-alkylamino, acylamino, carboxy, amido, mono- and di-alkylamido, alkylthio, 10 alkylsulfinyl, and alkylsulfonyl.

R<sub>4</sub> is hydrogen, alkyl or aralkyl; and

R<sub>5</sub>, R<sub>6</sub>, R<sub>7</sub> and R<sub>8</sub> are independently hydrogen, alkoxy or aralkoxy; or 15 a pharmaceutically acceptable salt thereof.

#### Detailed Description and Preferred Embodiments

As employed above and throughout this disclosure, the following terms, 20 unless otherwise indicated, shall be understood to have the following meanings:

"Monocyclic aryl" or "monocyclic heteroaryl" means a carbocyclic or heterocyclic aromatic ring. Preferred rings include phenyl, pyrrolyl, thienyl, furyl, thiazolyl, imidazolyl, pyrazolyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, 25 isothiazolyl, isoxazolyl and oxazolyl.

"Bicyclic aryl" or "bicyclic heteroaryl" means a bicyclic ring system composed of two fused rings at least one of which is a carbocyclic or heterocyclic aromatic ring. Preferred rings include substituted and unsubstituted 30 naphthyl, tetralinyl, 1,2,3,4-tetrahydroquinolyl, benzofuryl, benzothienyl, indolyl, indolinyl, 1,3-benzodioxolyl, benzodioxanyl, quinolinyl, tetrahydroquinolinyl, isoquinolinyl, quinazolinyl and quinoxalinyl.

"Monocyclic cycloalkyl" means a cyclic aliphatic ring comprising from 35 about four to about seven carbon atoms such as substituted and unsubstituted cyclopentyl, cyclohexyl and cycloheptyl.

"Bicyclic cycloalkyl" means a saturated bicyclic ring system composed of two fused aliphatic rings comprising about 8 to about 12 carbon atoms such as substituted and unsubstituted decalinyl.

5        "Monocyclic heterocycloalkyl" means a saturated cyclic aliphatic ring comprising from about four to about seven atoms and includes 1-2 hetero atoms selected from N, O and S provided said hetero atoms are not vicinal oxygen and/or sulfur atoms such as substituted and unsubstituted piperidinyl, piperazinyl and morpholinyl.

10        "Bicyclic heterocycloalkyl" means a saturated bicyclic ring system comprising two fused aliphatic rings of about 8 to about 12 atoms and includes 1-3 hetero atoms selected from N, O and S provided said hetero atoms are not vicinal oxygen and/or sulfur atoms such as substituted and unsubstituted  
15        decahydroquinolinyl.

      "Alkyl" means a saturated aliphatic hydrocarbon, either branched- or straight-chained. Preferred alkyl is "loweralkyl" having about 1 to about 6 carbon atoms. Examples of alkyl include methyl, ethyl, n-propyl, isopropyl, butyl, sec-butyl, t-butyl, amyl and hexyl.  
20

      "Alkoxy" refers to an alkyl-O-group. Preferred alkoxy groups include methoxy, ethoxy, propoxy and butoxy.

25        "Aryloxy" refers to an aryl-O-group. The preferred aryloxy group is phenoxy.

      "Aralkyl" means an alkyl group substituted by an aryl radical. The preferred aralkyl groups are benzyl or phenethyl.  
30

      The preferred aralkoxy groups are benzyloxy and phenethoxy.

      The preferred acyloxy groups are acetoxy and benzyloxy;

35        "Halo" means halogen. Preferred halogens include chloride, bromide and fluoride.

The preferred haloalkyl groups are mono-, di- and trifluoromethyl.

More specifically, the preferred A monocyclic aryl or heteroaryl rings include substituted or unsubstituted phenyl, pyrrolyl, thienyl, furyl, thiazolyl, imidazolyl, pyrazolyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, isothiazolyl, isoxazolyl and oxazolyl.

The preferred A bicyclic aryl or heteroaryl rings include substituted and unsubstituted naphthyl, tetralinyl, 1,2,3,4-tetrahydroquinolinyl, benzofuryl, benzothienyl, indolyl, indolinyl, 1,3-benzodioxolyl, benzodioxanyl, quinolinyl, tetrahydroquinolinyl, isoquinolinyl, quinazolinyl and quinoxalinyl.

The preferred A monocyclic cycloalkyl or heterocycloalkyl rings include substituted and unsubstituted cyclopentyl, cyclohexyl, cycloheptyl, decalanyl, piperdinyl, piperazinyl, morpholinyl or decahydroquinolinyl.

The preferred A bicyclic cycloalkyl or heterocycloalkyl rings include substituted and unsubstituted decalanyl or decahydroquinolinyl.

The more preferred compounds are those where:

A is substituted and unsubstituted phenyl, pyridyl, thienyl, furyl, pyrazolyl, naphthyl, tetralinyl, 1,2,3,4-tetrahydroquinolinyl, indolyl, indolinyl, quinolinyl, tetrahydroquinolinyl, cyclohexyl, piperdinyl or piperazinyl;

X is a bond, O, S, or NR<sub>4</sub>;

R is hydrogen, alkyl, alkoxy, halo, haloalkyl, alkylthio, alkylsulfinyl, alkylsulfonyl, phenyl and aralkyl;

R<sub>4</sub> is hydrogen, alkyl or aralkyl; and

R<sub>5</sub>, R<sub>6</sub>, R<sub>7</sub>, and R<sub>8</sub> are independently hydrogen or alkoxy.

The most preferred compounds are those where:

A is substituted and unsubstituted phenyl, naphthyl or indolyl;

X is a bond;

5 R is hydrogen, methoxy, ethoxy, chloro, trifluoromethyl, methylsulfonyl, phenyl and benzyl;

R<sub>4</sub> is hydrogen, methyl or benzyl; and

10 R<sub>5</sub>, R<sub>6</sub>, R<sub>7</sub>, and R<sub>8</sub> are independently hydrogen or methoxy.

Compounds within the scope of this invention are selective tyrosine kinase inhibitors of the human epidermal growth factor receptor type 2 (HER2). It is believed that therapeutically useful PTK inhibiting compounds have a specific tyrosine kinase activity which is capable of catalyzing  
15 autophosphorylation. In addition these compounds should inhibit growth factor-induced cell proliferation. Compounds meeting these criteria are of considerable value and are particularly useful in the practice of the present invention. Compounds exhibiting selectivity for the above receptor are described herein.

20 The compounds of this invention may be useful in the form of the free base, in the form of salts and as a hydrate. All forms are within the scope of the invention. Acid addition salts may be formed and are simply a more convenient form for use; and in practice, use of the salt form inherently amounts to use of the base form. The acids which can be used to prepare the acid addition salts  
25 include preferably those which produce, when combined with the free base, pharmaceutically acceptable salts, that is, salts whose anions are non-toxic to the animal organism in pharmaceutical doses of the salts, so that the beneficial properties inherent in the free base are not vitiated by side effects ascribable to the anions. Although pharmaceutically acceptable salts of said basic compound  
30 are preferred, all acid addition salts are useful as sources of the free base form even if the particular salt per se is desired only as an intermediate product as, for example, when the salt is formed only for purposes of purification and identification, or when it is used as an intermediate in preparing a pharmaceutically acceptable salt by ion exchange procedures.

35 Pharmaceutically acceptable salts within the scope of the invention include those derived from the following acids: mineral acids such as

hydrochloric acid, sulfuric acid, phosphoric acid and sulfamic acid; and organic acids such as acetic acid, citric acid, lactic acid, tartaric acid, malonic acid, methanesulfonic acid, ethanesulfonic acid, benzenesulfonic acid, p-toluenesulfonic acid, cyclohexylsulfamic acid, quinic acid, and the like.

5

The corresponding acid addition salts comprise the following: hydrochloride, sulfate, phosphate, sulfamate, acetate, citrate, lactate, tartrate, methanesulfonate, ethanesulfonate, benzenesulfonate, p-toluenesulfonate, cyclohexylsulfamate and quinate, respectively.

10

The acid addition salts of the compounds of this invention are prepared either by dissolving the free base in aqueous or aqueous-alcohol solution or other suitable solvents containing the appropriate acid and isolating the salt by evaporating the solution, or by reacting the free base and acid in an organic solvent, in which case the salt separates directly or can be obtained by concentration of the solution.


15

The compounds of this invention may be prepared by employing procedures known in the literature starting from known compounds or readily prepared intermediates. Exemplary general procedures follow.

20


In general the compounds useful for the method for the selective treatment of cell growth and differentiation characterized by activity of the human epidermal growth factor receptor type 2 may be prepared by the coupling reaction of a palladium catalyzed aryl or heteroaryl stannane with an aryl or heteroaryl halide or triflate.

25




where Y is halogen or triflate and Z is trialkylstannane and R is as previously described.


- 5 The 4-haloquinazoline starting materials are prepared in the classical way using anthranilic acid derivatives and formamide at reflux to provide the intermediate quinazolinones. Subsequent treatment with  $\text{POCl}_3$  at about  $110^\circ\text{C}$  for about two hours provides the chloroquinazolines. The final products are prepared via a condensation with the appropriate aniline derivative in a polar solvent such as ethanol. In the case of the phenoxy or thiophenoxy derivatives,
- 10 the metal salt (preferably Na) is prepared and refluxed for several hours with the appropriate haloquinazoline in a solvent such as THF.



- 15 The aryl and heteroarylstannanes may be prepared from the corresponding halide (preferably bromide or iodide) by conversion to the aryllithium by reaction with t-butyllithium at decreased temperatures, preferably about  $-78^\circ\text{C}$  followed by reaction with a halotrialkylstannane.



Of course these products may also be prepared in the reverse manner using the aryl or heteroarylhalides with the the corresponding stannane.



5

The quinazoline stannanes intermediates may be prepared by the action of trimethyltin sodium on aryl halides as described in Chem. Pharm. Bull. 1982, 30, 1731-1737:

- 10 The preparation of the compounds useful in this invention are described in Applicants' copending applications PCT/US94/14180 having International filing date December 8, 1994 and designating the United States as a contracting state, U.S. Serial No. 08/166,199, filed December 10, 1993 and U.S. Serial No. 08/229,886, filed April 19, 1994 all of which this application claims priority.
- 15 PCT/US94/14180, U.S. Serial No. 08/166,199 and U.S. Serial No. 08/229,886 are hereby incorporated herein by reference.

Further, the following examples are representative of the processes used to synthesis the compounds of this invention.

20

The below examples and those described in PCT/US94/14180 and U.S. Serial No. 08/166,199 may be followed to prepare any of the desired compounds of this invention. A representative list of compounds which may be prepared is shown below.

25

#### EXAMPLE 1

##### 4-(3-chlorophenoxy)-6,7-dimethoxyquinazoline

30

THF (5 ml) and NaH (60% disp in oil, approx. 28 mg) is added to a dry flask maintained under inert atmosphere at room temperature. 3-Chlorophenol (0.09 g) is added as a soln. in THF (1 mL) and stirring is continued until the solution became clear. 4-Chloro-6,7-dimethoxyquinazoline is added all at once

(as the solid) and stirring was maintained overnight at RT. The solution is partitioned between CH<sub>2</sub>Cl<sub>2</sub> and 5% NaOH. The organic layer is washed with brine, dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated. Flash column chromatography (40% EtOAc/Hex) provided the pure compound. An analytical sample is obtained by  
5 recrystallization from EtOAc/Hex to provide 4-(3-chlorophenoxy)-6,7-dimethoxyquinazoline (0.05 g), white needles, m.p. 152-153°C.

## EXAMPLE 2

### 4-(1-methylsulphonylindol-3-yl)-6,7-dimethoxyquinazoline

#### 10 Step A      N-methylsulfonyl-3-trimethylstannylindole

A solution of 5 g (15.57 mmol) of N-methylsulfonyl-3-iodoindole (5.1 g; 15.57 mmol) of hexamethylditin and 0.89 g (0.78 mmol) of Pd (PPh<sub>3</sub>)<sub>4</sub> in 75 mL of dry toluene is flushed thoroughly with nitrogen and heated to 90°C for 4 hours. The mixture is then evaporated and chromatographed on silica gel  
15 (eluting with hexane and then with 10% ethyl acetate/hexane to give N-methylsulfonyl-3-trimethylstannylindole which is used directly in the next step.

#### Step B      4-(1-methylsulphonylindol-3-yl)-6,7-dimethoxyquinazoline

A solution of 1.33 g (4.01 mmol) of N-methylsulfonyl-3-trimethylstannylindole, 750 mg (3.34 mmol) of 4-chloro-6,7-dimethoxyquinazoline and  
20 0.19 g (5 mol % 0.16 mmol) of Pd (PPh<sub>3</sub>)<sub>4</sub> in 10 ml of dry dimethylformamide is flushed thoroughly with nitrogen and heated to 90°C for 12 hours. The reaction mixture is diluted with methylene chloride washed with 10% ammonium hydroxide and stirred vigorously and then washed with water and the combined organics are washed with brine (75 mL), dried (MgSO<sub>4</sub>) and evaporated to  
25 dryness. Recrystallization from ethyl acetate yields 4-(1-methylsulphonylindol-3-yl)-6,7-dimethoxyquinazoline (m.p. >220°C).

The above examples may be followed to prepare any of the desired  
30 compounds of this invention. A representative list of compounds which may be prepared are shown below.

6,7-dimethoxy-4-naphthalen-2-ylethynylquinazoline, m.p. 158-161°C

35 4-(4-hydroxyphenyl)-6,7-dimethoxyquinazolinehydrochloride,

- m.p. >270°C (dec)
- 4-(naphthalen-1-yl)-6,7-dimethoxyquinazoline, m.p. 144-147°C
- 5 4-(naphthalen-2-yl)-6,7-dimethoxyquinazoline, m.p. 115-118°C
- 4-phenylacetylenyl-6,7-dimethoxyquinazoline, m.p. 146-148°C
- 4-(3-fluoro-4-methoxyphenyl)-6,7-dimethoxyquinazoline, m.p. 207-210°C
- 10 4-(3-phenylphenyl)-6,7-dimethoxyquinazoline, m.p. 160-163°C
- 4-(2-phenylethylenyl)-6,7-dimethoxyquinazoline, m.p. 168-169°C
- 4-(2-methoxypyridin-5-yl)-6,7-dimethoxyquinazoline, m.p. 175-176°C
- 15 4-(1-benzylindol-3-yl)-6,7-dimethoxyquinazoline, m.p. 148-150°C
- 4-(indol-3-yl)-6,7-dimethoxyquinazoline, m.p. >240°C (dec)
- 20 4-(1-methylindol-3-yl)-6,7-dimethoxyquinazoline hydrochloride,  
m.p. >230°C (dec)
- 4-(1-methylsulphonylindol-3-yl)-6,7-dimethoxyquinazoline, m.p. >220°C (dec)
- 25 4-(4-phenylpiperidin-1-yl)-6,7-dimethoxyquinazoline, m.p. 150-151°C
- 4-[4-(3-chlorophenyl)piperazin-1-yl]-6,7-dimethoxyquinazoline, m.p. 155-156°C
- 30 4-(N-methyl-3,4,5-trimethoxyanilino)-6,7-dimethoxyquinazoline,  
m.p. 149-151°C
- (+ -)-4-(2-methyl-1,2,3,4-tetrahydroquinolin-1-yl)-6,7-dimethoxyquinazoline  
hydrochloride, m.p. 198-201°C (dec)
- 35 4-(1,2,3,4-tetrahydroquinolin-1-yl)-6,7-dimethoxyquinazoline hydrochloride, m.p.  
195-197°C (dec)

- 4-(5,6,7,8-tetrahydronaphthalen-1-yl)amino-6,7-dimethoxyquinazoline  
hydrochloride, m.p. 219-222°C
- 5 4-(3,6-dioxananilino)-6,7-dimethoxyquinazoline, m.p. 267-269°C (dec)
- 4-phenylacetylenyl-6,7-dimethoxyquinazoline, m.p. 146-148°C
- 4-(indol-1-yl)-6,7-dimethoxyquinazoline, m.p. 166-167°C
- 10 4-(N-methyl-4-methoxyanilino)-6,7-dimethoxyquinazoline hydrochloride,  
m.p. 202-205°C
- 4-(N-methyl-4-chloroanilino)-6,7-dimethoxyquinazoline hydrochloride,  
m.p. 220-222°C
- 15 4-(2,3-dihydroindol-1-yl)-6,7-dimethoxyquinazoline hydrochloride,  
m.p. 226-229°C (dec)
- 20 N-(6,7-dimethoxyquinazolin-4-yl)-N-methyl-N-(3-trifluoromethylphenyl)amine  
hydrochloride, m.p. 240-243°C
- N-(3-chlorophenyl)-N-(6,7-dimethoxyquinazolin-4-yl)-N-methylamine  
hydrochloride, m.p. 235-237°C
- 25 N-(3-chlorophenyl)-N-(quinazolin-4-yl)-N-methylamine hydrochloride,  
m.p. 233-235°C
- 6,7-dimethoxy-4-naphthalen-1-yl-ethynylquinazoline, m.p. 175-177°C
- 30 4-(thien-3-yl)-6,7-dimethoxyquinazoline, m.p. 148.5-151.5°C
- 4-benzyl-6,7-dimethoxyquinazoline, m.p. 122.5-125°C
- 35 (6,7-dimethoxyquinazolin-4-yl)-5-indazolylamine hydrochloride,  
m.p. 261-263°C (dec)

N-(6,7-dimethoxyquinazolin-4-yl)-N-phenyl-N-ethylamine hydrochloride,  
m.p. 227-230°C (dec)

5 N-benzyl-N-(6,7-dimethoxyquinazolin-4-yl)-N-phenylamine hydrochloride,  
m.p. 269-271°C

N-(6-chloroquinazolin-4-yl)-N-methyl-N-phenylamine, m.p. 106-108°C

10 N-(3-chloro-phenyl)-N-(6,7-dimethoxyquinazolin-4-yl)-N-ethylamine  
hydrochloride, m.p. 261-263°C

N-(6,7-dimethoxyquinazolin-4-yl)-N-methyl-N-p-tolylamine hydrochloride,  
m.p. 230-234°C (dec)

15 N-benzyl-N-(6,7-dimethoxyquinazolin-4-yl)amine, m.p. 220-225°C

N-(4-methoxybenzyl)-N-(6,7-dimethoxyquinazolin-4-yl)amine, m.p. 194-198°C

20 N-(3,5-dimethoxybenzyl)-N-(6,7-dimethoxyquinazolin-4-yl)amine hydrochloride,  
m.p. 265-269°C

4-(3,4,5-trimethoxyphenoxy)-6,7-dimethoxyquinazoline, m.p. 228-232°C

25 N-(quinazolin-4-yl)-N-phenyl-N-methylamine hydrochloride, m.p. 242-246°C (dec)

N-(6,7-dimethoxyquinazolin-4-yl)-N-(4-morpholin-4-ylphenyl)amine hydrochloride,  
m.p. 231-235°C (dec)

30 4-(3-methoxythiophenoxy)-6,7-dimethoxyquinazoline, m.p. 139.5-141.5°C

4-[N-(5-indanyl)amino]-6,7-dimethoxyquinazoline hydrochloride,  
m.p. 244-246°C (dec)

4-(3-chlorothiophenoxy)-6,7-dimethoxyquinazoline, m.p. 152-153.5°C

- 4-(3-aminopyrazolyl)-6,7-dimethoxyquinazoline hydrochloride,  
m.p. 262-264°C (dec)
- 5 4-(1,4-benzodioxan-6-ylamino)-6,7-dimethoxyquinazoline hydrochloride,  
m.p. 267-269°C (dec)
- 6,7-dimethoxy-4-( $\alpha$ -naphthylamino)quinazoline hydrochloride, m.p. >250°C
- 10 6,7-dimethoxy-4-( $\beta$ -naphthylamino)quinazoline hydrochloride, m.p. >250°C
- 4-(cyclohexylanilino)-6,7-dimethoxyquinazoline, m.p. 239-244°C
- 4-(3,4,5-trimethoxyanilino)-6,7-dimethoxyquinazoline hydrochloride,  
m.p. 260-265°C
- 15 6,7-dimethoxy-4-(N-methylanilino)quinazoline hydrochloride, m.p. >230°C
- 4-(3-chlorophenoxy)-6,7-dimethoxyquinazoline, m.p. 152-153°C
- 20 6,7-dimethoxy-4-(1-naphthylthio)-quinazoline, m. p. 174.5-176.5°C
- 6,7-dimethoxy-4-(2-naphthylthio)-quinazoline, m. p. 178-179°C
- 25 6,7-dimethoxy-4-(1-naphthyloxy)-quinazoline, m. p. 214-215.5°C
- 6,7-dimethoxy-4-(2-naphthyloxy)-quinazoline, m. p. 169-170°C
- N-(6,7-dimethoxy-quinolazolin-4-yl)-N-(naphth-2-yl)-N-ethylamine hydrochloride,  
m. p. 236-239°C (dec)
- 30 6,7-dimethoxy-4-(naphthalene-2-sulfinyl)quinazoline, m. p. 182.5-185°C
- 6,7-dimethoxy-4-(naphthalene-2-sulfonyl)quinazoline
- 35 4-(3-chloroanilino)-6,7-dimethylquinazoline hydrochloride, m.p. 271-274°C

4-(3,5-dimethylanilino)-6,7-dimethylquinazoline hydrochloride,  
m.p. >275°C

5 4-(N-methyl-4-methylanilino)-6,7-dimethylquinazoline hydrochloride,  
m.p. 235-238°C

6,7-dimethyl-4-(1-naphthylamino)quinazoline hydrochloride,  
m.p. 244-247°C

10 6,7-dimethyl-4-(7-trifluoromethyl-3,4-dihydro-2H-quinolin-1-yl)quinazoline  
hydrochloride, m.p. 240°C

4-(N-methyl-3-methylanilino)-6,7-dimethylquinazoline hydrochloride,  
m.p. 205-207°C

15

4-(3-chlorophenylthio)-6,7-dimethylquinazoline hydrochloride,  
m.p. 197-202°C

20 4-(1-naphthylthio)-6,7-dimethylquinazoline hydrochloride,  
m.p. 204-209°C

4-(3,4-dimethoxyphenylthio)quinazoline, m.p. 115-117°C

25

Preparation of Pharmaceutical Compositions and  
Pharmacological Test Section

30 To determine the effectiveness of compounds of this invention, the  
pharmacological tests described below, which are accepted in the art and  
recognized to correlate with pharmacological activity in mammals, are utilized.  
Compounds within the scope of this invention have been subjected to these  
various tests, and the results obtained are believed to correlate to useful  
inhibition of growth factor-induced cell proliferation. The below described tests  
are useful in determining the inhibition of the compounds of this invention.

### Experimental Procedures

Materials: The test material was dissolved extemporaneously in DMSO to make up stock solutions which were subsequently diluted in culture medium to reach the desired compound concentrations (vehicle final concentration of 0.1%). The materials used are as follows: Dulbecco's Modified Eagle Medium (DMEM), RPMI 1640, fetal calf serum (FCS), and Penicillin-Streptomycin solution (10,000 IU/ml penicillin and 10,000 µg/ml streptomycin), were from GIBCO BRL. 3-(4, 5-dimethylthiazol-2 yl)-2, 5-diphenyltetrazolium bromide (MTT), and geneticin were from Sigma. Radiolabeled [ $\gamma$ -<sup>32</sup>P]ATP (cat#NEG002H; 3,000 Ci/mmol), was purchased from NEN. Anti-c neu antibodies (Ab-5; cat#OP39) were from Oncogene Science. Mouse monoclonal anti-phosphotyrosine antibodies (monoclonal IgG2bk) were from UBI and horseradish peroxidase-conjugated rabbit anti-mouse IgG were from Nordic Immunological Laboratories. Stromelysin-CAT plasmid DNA construct, consisting of the chloramphenicol acetyl transferase cDNA under the control of the stromelysin promoter fragment -1100/+8 was prepared by the procedure of C. Wasylyk, A. Gutman, R. Nicholson and B. Wasylyk, (1991) *EMBO J.*, **10**: A127-A134. All other chemicals were of the best quality available.

Cell lines and cell culture: Mouse embryo fibroblasts (NIH3T3), human epidermoid (A431) carcinoma cell line, and breast (SK-BR-3) and ovary (SK-OV-3) adenocarcinoma cell lines, human lung (A549), and breast (BT474 and BT20) carcinoma cell lines were purchased from the ATCC. NIH3T3 cells transfected with HER2 construct and overexpressing p185HER2, NIH/HER2 were prepared according to R. M. Hudziak, J. Schlessinger and A. Ullrich (1987) *Proc. Natl. Acad. Sci. USA*, **84**: 7159-7163. NIH/Ha-Ras cell line was established by transforming NIH3T3 cells with Val2Ha-ras according to I. Barlat, F. Schweighoffer, M. C. Chevalier-Multon, M. Duchesne, I. Faith, D. Landais, M. Jacquet and B. Tocqué (1993) *Oncogene*, **8**: 215-218. NIH3T3 cells transfected with activated v-src containing mutation Y527F (NIH/v-Src) were obtained from Dr. B. Wasylyk. HIR3.5 (NIH/IR) cell line represents NIH3T3 cells overexpressing the human insulin receptor and was provided by Dr. J. Whitaker. All cell lines were cultured at 37°C in a CO<sub>2</sub> incubator (5% CO<sub>2</sub>/95% humidified air atmosphere) in medium supplemented with 10% FCS and 1% penicillin/streptomycin solution. Unless specified in the text, the culture medium was changed every other day.

In vitro p185<sup>HER2</sup> autophosphorylation assay: ER22 or A431, and NIH/HER2 cell lines were used as sources for p185<sup>Her2</sup>. Subconfluent cell monolayers were lysed at 4°C for 10 min in HNEG buffer (HNEG: 50mM Hepes buffer, pH 7.5, 150mM NaCl, 1 mM EDTA and 10% Glycerol) containing 1% Triton X-100 and 1mM PMSF. Cell lysates were then diluted with HNEG buffer containing 0.1% Triton X-100 and 1% BSA (lysis buffer), and cell extracts were clarified by centrifugation at 12,000.g for 5 min. Autophosphorylation assays were preformed as described by S. M. Smyth, I. Stefanova, F. Hartman, I.D. Horak, N. Osherov, A. Levitzki, and T. R. Burke (1993) *J. Med. Chem.*, **36**: 3010-3014. Briefly, 96 well U-bottom plastic plates were coated at 37°C for 2 h with 100µl of goat anti-mouse immunoglobulin (Biosys) at a concentration of 10µg/ml. After several washes with PBS containing 0.05% Tween, 100 µl of anti-c-neu antibodies were incubated for 2 h at 37°C (both antibodies were used at 1µg/ml). Unoccupied binding sites were blocked by incubation for 1 h at 37°C with 2% BSA in PBS. Cell lysates were incubated in coated wells for 1 h at 4°C. After several washes with lysis buffer autophosphorylation assay was performed directly in the wells in 25mM Hepes buffer, pH 7.4, containing 2mM MnCl<sub>2</sub>, 0.1% Triton and 5 µCi [ $\gamma$ -<sup>32</sup>P]ATP for 20 min at room temperature in the absense or the presence of Applicants' compounds. Phosphorylation reaction was quenched by adding Laemmli's sample buffer prepared according to U. K. Laemmli (1970) (*Nature*, **225**: 680-685 and [<sup>32</sup>P]-labeled receptors were analyzed by SDS-PAGE on 4-12% polyacrylamide gradient gels. Phosphorylation intensities were estimated by scanning the dried gels on an Instant Imager (Packard).

25

Intact cell tyrosine phosphorylation: Content cell monolayers were serum-starved overnight after which they were incubated with the indicated compound concentrations for 2 h. Medium was then aspirated and cells were quenched by adding Laemmli's SDS-sample buffer directly on cell monolayers. Samples were then treated at 100°C for 5 min before being tested for phosphotyrosine-containing proteins. Proteins were fractionated by SDS-Page on 4-20% polyacrylamide gradient gels, after which proteins were electrophoretically transfered to polyvinylidene difluoride membranes (PVDF membrane, PolyScreen, NEN). Immunological detection of phosphotyrosine-containing proteins was performed using a mouse monoclonal anti-phosphotyrosine antibody. Blots were developed by the Enhanced ChemiLuminescence method (ECL,NEN) employing horseradish peroxidaseconjugated rabbit anti-mouse IgG.

Cell proliferation: Cells were seeded onto 24 well cell culture plates at about 20,000 cells per well. Cells were allowed to adhere to plastic for 8 h in 1 ml culture medium, after which cells were cultured in the presence of various concentrations of compounds for 72-96 h. Following the indicated incubation times, cell number per well was estimated. As an assay of relative viable cell number, mitochondrial reduction of MTT was used following the procedure of M. C. Alley, D. A. Scudiero, A. Monks, M.L. Hursey, M. J. Czerwinski, D. L. Fine, B. J. Abbot, J. G. Mayo, R. H. Shoemaker and M. R. Boyd (1988) *Cancer Res.* 48: 589-601. Briefly, 100  $\mu$ l of a 5 mg/ml solution of MTT in phosphate-buffered saline was added to each well, and plates were incubated for 4 h at 37°C in a CO<sub>2</sub> incubator. 650  $\mu$ l of medium was then removed and replaced by 750  $\mu$ l of an isopropyl alcohol/HCl (1N) solution (25:1) in order to dissolve the dark purple crystals of formazan formed in the mitochondria of living cells. After incubation for 5-10 min at room temperature under agitation, 200- $\mu$ l aliquots from each well were transferred to 96-well cell culture plates, and since the extent of obtained bluish color was directly proportional to cell number, this was estimated by spectrophotometry at 590 nm in a microplate autoreader.

Anchorage-independent cell growth: Anchorage-independent cell growth was investigated by examining the colony-forming capability of the considered cells suspended in soft-agar. Experiments were performed using 50-mm diameter cell culture dishes. A 4-ml cell-free feeder underlayer consisted of 0.5% agar in medium supplemented with 10% FCS and the indicated concentrations of compounds. The 4-ml overlayer contains about 10,000 cells in 0.3% agar in medium supplemented with 10% FCS and the corresponding concentrations of compounds. After incubation for 2 weeks at 37°C, the number of colonies was determined.






Cell transfection and CAT assay: Stromelysin-CAT plasmid DNA construct (STRM-CAT) was introduced into cells by transfection using the lipofectAMINE reagent. Briefly, 50-70% confluent culture dishes (3.5-cm) of cells were exposed to 1  $\mu$ g of plasmid DNA and 10  $\mu$ g of lipofectAMINE reagent in 1 ml of serum-free DMEM for 4 h at 37°C. Cells were then incubated for 36 h at 37°C with DMEM supplemented with 0.5% FCS and the indicated amounts of compounds. Cells were detached off the plates by incubation with PBS containing 3 mM EDTA and pelleted by centrifugation for 5 min at 1,500 rpm. Cells were resuspended in 0.25 M Tris/HCl buffer, pH 7.8 and were subjected

to repeated freeze-thawing cycles. Cell extracts were heated to 65°C for 15 min and, after cooling, were microfuged for 15 min at 14,000 rpm. Supernatants were assayed for CAT activity following the procedure of J. R. Neumann, C. A. Morency and K. O. Russian (1987) *BioTechniques*, 5: 444-447.

5

Compounds within the scope of this invention exhibit significant specific activity as protein tyrosine kinase inhibitors and possess therapeutic value for inhibiting growth factor-induced cell proliferation. Further, compounds of this invention are specific inhibitors of the human epidermal growth factor receptor type 2 and are therefore useful for treating cell growth and differentiation.

The following table shows examples of representative compounds of this invention and their test results as determined by the above inhibition of the human epidermal growth factor receptor type 2. The results obtained by the above experimental methods evidence the useful HER2 receptor protein tyrosine kinase inhibition properties of compounds within the scope of the present invention and possess therapeutic value in the regulation of abnormal cell growth.

	X	$R_5$	$R_6$	$R_7$	$R_8$	HER-2
	bond	H	OCH <sub>3</sub>	OCH <sub>3</sub>	H	20% at 5μM
	bond	H	OCH <sub>3</sub>	OCH <sub>3</sub>	H	10% at 5μM
	bond	H	OCH <sub>3</sub>	OCH <sub>3</sub>	H	0.5-1μM
	bond	H	OCH <sub>3</sub>	OCH <sub>3</sub>	H	10% at 5μM

20

The compounds of the present invention can be administered to a mammalian host in a variety of forms adapted to the chosen route of administration, i.e., orally, or parenterally. Parenteral administration in this respect includes administration by the following routes: intravenous,  
5 intramuscular, subcutaneous, intraocular, intrasynovial, transepithelial including transdermal, ophthalmic, sublingual and buccal; topically including ophthalmic, dermal, ocular, rectal and nasal inhalation via insufflation and aerosol and rectal systemic.

10 The active compound may be orally administered, for example, with an inert diluent or with an assimilable edible carrier, or it may be enclosed in hard or soft shell gelatin capsules, or it may be compressed into tablets, or it may be incorporated directly with the food of the diet. For oral therapeutic administration, the active compound may be incorporated with excipient and used in the form of  
15 ingestible tablets, buccal tablets, troches, capsules, elixirs, suspensions, syrups, wafers, and the like. Such compositions and preparations should contain at least 0.1% of active compound. The percentage of the compositions and preparations may, of course, be varied and may conveniently be between about 2 to about 6% of the weight of the unit. The amount of active compound in such  
20 therapeutically useful compositions is such that a suitable dosage will be obtained. Preferred compositions or preparations according to the present invention are prepared so that an oral dosage unit form contains between about 1 and 1000 mg of active compound.

25 The tablets, troches, pills, capsules and the like may also contain the following: A binder such as gum tragacanth, acacia, corn starch or gelatin; excipients such as dicalcium phosphate; a disintegrating agent such as corn starch, potato starch, alginic acid and the like; a lubricant such as magnesium stearate; and a sweetening agent such as sucrose, lactose or saccharin may be  
30 added or a flavoring agent such as peppermint, oil of wintergreen, or cherry flavoring. When the dosage unit form is a capsule, it may contain, in addition to materials of the above type, a liquid carrier. Various other materials may be present as coatings or to otherwise modify the physical form of the dosage unit. For instance, tablets, pills, or capsules may be coated with shellac, sugar or  
35 both. A syrup or elixir may contain the active compound, sucrose as a sweetening agent, methyl and propylparabens as preservatives, a dye and flavoring such as cherry or orange flavor. Of course, any material used in

preparing any dosage unit form should be pharmaceutically pure and substantially non-toxic in the amounts employed. In addition, the active compound may be incorporated into sustained-release preparations and formulations.

5

The active compound may also be administered parenterally or intraperitoneally. Solutions of the active compound as a free base or pharmacologically acceptable salt can be prepared in water suitably mixed with a surfactant such as hydroxypropylcellulose. Dispersion can also be prepared in  
10 glycerol, liquid polyethylene glycols, and mixtures thereof and in oils. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms.

The pharmaceutical forms suitable for injectable use include sterile  
15 aqueous solutions or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. In all cases the for must be sterile and must be fluid to the extent that easy syringability exists. It may be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi.  
20 The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), suitable mixtures thereof, and vegetable oils. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of  
25 dispersion and by the use of surfactants. The prevention of the action of microorganisms can be brought about by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars or sodium chloride. Prolonged absorption of the injectable  
30 compositions can be brought about by use of agents delaying absorption, for example, aluminum monostearate and gelatin.

Sterile injectable solutions are prepared by incorporating the active compound in the required amount in the appropriate solvent with various of the  
35 other ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the various sterilized active ingredient into a sterile vehicle which contains the basic


dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and the freeze drying technique which yield a powder of the active ingredient plus any  
5 additional desired ingredient from previously sterile-filtered solution thereof.

The therapeutic compounds of this invention may be administered to a mammal alone or in combination with pharmaceutically acceptable carriers, as noted above, the proportion of which is determined by the solubility and  
10 chemical nature of the compound, chosen route of administration and standard pharmaceutical practice.

The dosage of the present therapeutic agents which will be most suitable for prophylaxis or treatment will vary with the form of administration, the  
15 particular compound chosen and the physiological characteristics of the particular patient under treatment. Generally, small dosages will be used initially and if necessary, will be increased by small increments until the optimum effect under the circumstances is reached. The therapeutic human dosage, based on physiological studies using rats, will generally be from about 0.01 mg to about 100  
20 mg/kg of body weight per day or from about 0.4 mg to about 10 g or higher although it may be administered in several different dosage units from once to several times a day. Oral administration requires higher dosages.

## CLAIMS

- 5 1. A method for the selective treatment of cell growth and differentiation characterized by activity of the human epidermal growth factor receptor type 2 (HER2) comprising administering to a patient in need of such treatment an HER2 receptor inhibiting effective amount of a compound of the formula:



10

wherein

- A is a substituted or unsubstituted mono- or bi-cyclic aryl, heteroaryl, cycloalkyl or heterocycloalkyl ring system of about 5 to about 12 atoms and  
 15 where the substituents may be located at any appropriate position of the ring system and are described by R;

X is a bond, O, S, SO, SO<sub>2</sub>, OCH<sub>2</sub>, CR<sub>4</sub>=CR<sub>4</sub>, C≡C, NR<sub>4</sub> or NR<sub>4</sub>CH<sub>2</sub>;

20

R independently includes hydrogen, alkyl, phenyl, halophenyl, aralkyl, hydroxy, alkoxy, aryloxy, acyloxy, halo, haloalkyl, amino, mono- and di-alkylamino, acylamino, carboxy, amido, mono- and di-alkylamido, alkylthio, alkylsulfinyl, and alkylsulfonyl.

25

R<sub>4</sub> is hydrogen, alkyl or aralkyl; and

R<sub>5</sub>, R<sub>6</sub>, R<sub>7</sub> and R<sub>8</sub> are independently hydrogen, alkoxy or aralkoxy; or

a pharmaceutically acceptable salt thereof.

30

2. A method according to Claim 1 where:

A is a substituted or unsubstituted monocyclic aryl or heteroaryl ring selected from phenyl, pyrrolyl, thienyl, furyl, thiazolyl, imidazolyl, pyrazolyl,  
5 pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, isothiazolyl, isoxazolyl and oxazolyl.

3. A method according to Claim 1 where:

10 A is a substituted or unsubstituted bicyclic aryl or heteroaryl ring selected from naphthyl, tetralinyl, 1,2,3,4-tetrahydroquinoliny, benzofuryl, benzothienyl, indanyl, indolyl, indolinyl, 1,3-benzodioxolyl, benzodioxanyl, quinoliny, tetrahydroquinoliny, isoquinoliny, quinazoliny and quinoxaliny.

15

4. A method according to Claim 1 where:

A is a substituted or unsubstituted monocyclic cycloalkyl or heterocycloalkyl ring selected from cyclopentyl, cyclohexyl, cycloheptyl,  
20 decaliny, piperdiny, piperaziny, morpholiny or decahydroquinoliny.

5. A method according to Claim 1 where:

25 A is a substituted or unsubstituted bicyclic cycloalkyl or heterocycloalkyl ring selected from decaliny or decahydroquinoliny.

6. A method according to Claim 1 where:

30

A is substituted and unsubstituted phenyl, pyridyl, thienyl, furyl, pyrazolyl, naphthyl, tetralinyl, 1,2,3,4-tetrahydroquinoliny, indanyl, indolyl, indolinyl, quinoliny, tetrahydroquinoliny, cyclohexyl, piperdiny or piperaziny;

35

X is a bond, O, S, or NR<sub>4</sub>;

R is hydrogen, alkyl, alkoxy, halo, haloalkyl, alkylthio, alkylsulfinyl, alkylsulfonyl, phenyl and aralkyl;

R<sub>4</sub> is hydrogen, alkyl or aralkyl; and

5

R<sub>5</sub>, R<sub>6</sub>, R<sub>7</sub>, and R<sub>8</sub> are independently hydrogen or alkoxy.

7. A method according to Claim 6 where:

10

A is substituted and unsubstituted phenyl, naphthyl or indolyl;

X is a bond;

15

R is hydrogen, methoxy, ethoxy, chloro, trifluoromethyl, methylsulfonyl, phenyl and benzyl;

R<sub>4</sub> is hydrogen, methyl or benzyl; and

20

R<sub>5</sub>, R<sub>6</sub>, R<sub>7</sub>, and R<sub>8</sub> are independently hydrogen or methoxy.

8. A method according to Claim 2 where the compound administered is 4-(2-phenylethynyl)-6,7-dimethoxyquinazoline or a pharmaceutically acceptable thereof.

25

9. A method according to claim 3 where said compound administered is 6,7-dimethoxy-4-naphthalen-1-yl-ethynylquinazoline or a pharmaceutically acceptable thereof.

30

10. A method according to claim 6 where said compound administered is selected from:

35

4-(2-methoxypyridin-5-yl)-6,7-dimethoxyquinazoline,  
4-(4-phenylpiperidin-1-yl)-6,7-dimethoxyquinazoline,  
4-[4-(3-chlorophenyl)piperazin-1-yl]-6,7-dimethoxyquinazoline and

4-(1,2,3,4-tetrahydroquinolin-1-yl)-6,7-dimethoxyquinazoline  
or a pharmaceutically acceptable thereof.

5 11. A method according to claim 6 where said compound administered is  
selected from:

6,7-dimethoxy-4-( $\beta$ -naphthylamino)quinazoline,  
4-[N-(5-indanyl)amino]-6,7-dimethoxyquinazoline,  
N-benzyl-N-(6,7-dimethoxyquinazolin-4-yl)-N-phenylamine,  
10 6,7-dimethoxy-4-(N-methylanilino)quinazoline,  
N-(3-chlorophenyl)-N-(6,7-dimethoxyquinazolin-4-yl)-N-methylamine,  
4-(3-aminopyrazolyl)-6,7-dimethoxyquinazoline and  
4-(cyclohexylanilino)-6,7-dimethoxyquinazoline  
or a pharmaceutically acceptable thereof.

15

12. A method according to Claim 6 where the compound administered is 6,7-  
dimethoxy-4-( $\alpha$ -naphthylamino)quinazoline or a pharmaceutically acceptable salt  
thereof.

20

13. A method according to claim 7 where said compound administered is  
selected from:

4-(naphthalen-1-yl)-6,7-dimethoxyquinazoline,  
25 4-(naphthalen-2-yl)-6,7-dimethoxyquinazoline,  
4-(3-phenylphenyl)-6,7-dimethoxyquinazoline,  
4-(indol-3-yl)-6,7-dimethoxyquinazoline and  
4-(1-methylindol-3-yl)-6,7-dimethoxyquinazoline  
or a pharmaceutically acceptable salt thereof.


30

14. A method according to Claim 7 where the compound administered is  
4-(1-benzylindol-3-yl)-6,7-dimethoxyquinazoline or a pharmaceutically  
acceptable salt thereof.

35

15. A pharmaceutical composition for selectively treating cell growth and differentiation characterized by activity of the human epidermal growth factor receptor type 2 (HER2) comprising an HER2 receptor inhibiting effective amount of a compound of the formula:

5



wherein

10 A is a substituted or unsubstituted mono- or bi-cyclic aryl, heteroaryl, cycloalkyl or heterocycloalkyl ring system of about 5 to about 12 atoms and where the substituents may be located at any appropriate position of the ring system and are described by R;

15 X is a bond, O, S, SO, SO<sub>2</sub>, OCH<sub>2</sub>, CR<sub>4</sub>=CR<sub>4</sub>, C≡C, NR<sub>4</sub> or NR<sub>4</sub>CH<sub>2</sub>;

20 R independently includes hydrogen, alkyl, phenyl, halophenyl, aralkyl, hydroxy, alkoxy, aryloxy, acyloxy, halo, haloalkyl, amino, mono- and di-alkylamino, acylamino, carboxy, amido, mono- and di-alkylamido, alkylthio, alkylsulfinyl, and alkylsulfonyl.

R<sub>4</sub> is hydrogen, alkyl or aralkyl; and

R<sub>5</sub>, R<sub>6</sub>, R<sub>7</sub> and R<sub>8</sub> are independently hydrogen, alkoxy or aralkoxy; or

25 a pharmaceutically acceptable salt thereof, in admixture with a pharmaceutically acceptable carrier.

16. A composition according to Claim 15 where:

A is substituted and unsubstituted phenyl, pyridyl, thienyl, furyl, pyrazolyl, naphthyl, tetralinyl, 1,2,3,4-tetrahydroquinoliny, indanyl, indolyl, indoliny, quinoliny, tetrahydroquinoliny, cyclohexyl, piperdiny or piperaziny;

X is a bond, O, S, or NR<sub>4</sub>;

R is hydrogen, alkyl, alkoxy, halo, haloalkyl, alkylthio, alkylsulfinyl, alkylsulfonyl, phenyl and aralkyl;

R<sub>4</sub> is hydrogen, alkyl or aralkyl; and

R<sub>5</sub>, R<sub>6</sub>, R<sub>7</sub>, and R<sub>8</sub> are independently hydrogen or alkoxy.

17. A composition according to Claim 16 where the compound is 6,7-dimethoxy-4-( $\alpha$ -naphthylamino)quinazoline or a pharmaceutically acceptable salt thereof.

18. A composition according to Claim 16 where the compound is 4-(1-benzylindol-3-yl)-6,7-dimethoxyquinazoline or a pharmaceutically acceptable salt thereof.

19. 6,7-dimethoxy-4-( $\alpha$ -naphthylamino)quinazoline or a pharmaceutically acceptable salt thereof.

20. 4-(1-benzylindol-3-yl)-6,7-dimethoxyquinazoline or a pharmaceutically acceptable salt thereof.

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US96/09606

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : A61K 31/535, 31/495, 31/50, 31/505

US CL : 514/234.5, 249, 259

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbol(s))

U.S. : 514/234.5, 249, 259

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A, P	US 5,480,883 A (SPADA ET AL.) 02 January 1996 (02.01.96), see entire document.	1-20

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
* A document defining the general state of the art which is not considered to be of particular relevance		
* E earlier document published on or after the international filing date	X	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
* L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	Y	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
* O document referring to an oral disclosure, use, exhibition or other means		
* P document published prior to the international filing date but later than the priority date claimed	&	document member of the same patent family

Date of the actual completion of the international search

27 AUGUST 1996

Date of mailing of the international search report

17 SEP 1996

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

JEROME D. GOLDBERG

Telephone No. (703) 308-1235